Feeding Performance: How Modern Emulsifiers Boost Energy Use in Broilers

Dr. Attila Kovács DVM MVSc – Innovad Group



Modern poultry production focuses on obtaining maximum growth rates while maintaining economic sustainability as its core objective. The main production cost factor is feed, which requires efficient dietary energy utilisation to ensure profitability. The utilisation of dietary lipids—fats and oils—becomes essential within this framework. The ingredients serve as high-energy dense components, yet their efficiency depends on more than just their concentration in the diet. The efficiency depends on the digestive system's ability to properly break down and absorb them.

Lipid emulsification and bile salts – the reality.

Poultry lipid digestion begins in the small intestine, where bile salts act as primary amphipathic molecules, which the liver produces. These molecules possess both hydrophilic and hydrophobic domains, which allow them to encapsulate lipid droplets and convert them into more refined emulsions. The structural transformation of emulsions by these molecules makes larger surfaces available to pancreatic lipase, which performs triglyceride hydrolysis.

The digestion sequence consists of multiple coordinated steps, including emulsification followed by enzymatic breakdown to fatty

Continued on page 9 ▷

▷ *Continued from page 7*

acids and monoglycerides and bile salt-assisted micelle formation, which enables intestinal enterocyte uptake and re-esterification to triglycerides before packaging into chylomicrons for systemic circulation. This is a long and complex process; however, the efficiency of these digestion steps depends directly on the emulsification process.

Beyond physiology: functional ingredients to support lipid assimilation

The natural bile activity of broilers might not satisfy the needs of fast-growing birds or those consuming high-fat diets. The existing gap between natural fat digestion capabilities creates space for dietary emulsifying agent supplementation.

Lysolecithins derived from hydrolysed lecithins have emerged as significant emulsifying agents in the current reality. The process of enzymatic modification transforms natural phospholipids into molecules with superior surface properties. The substances enhance fat emulsification together with absorption while modifying cellular membrane properties to affect permeability and fluidity, and possibly signalling mechanisms connected to the immune system and metabolic responses.

Research indicates that particular lysolecithin compositions function as nutritional boosters and bioactive substances which affect both gut maturation and complete body physiological systems. The functional behaviour depends on various factors, including chemical structure as well as formulation methods and digestive characteristics that differ across species.

Engineering modern emulsifiers: the significance of hlb and bio-functional design

The effectiveness of an emulsifier in formulation science can be assessed through its hydrophilic-lipophilic balance (HLB), which indicates water versus oil molecule affinity. Emulsifiers with lower HLB values exhibit lipophilic properties, which stabilise water-in-oil emulsions, whereas high-HLB agents work best with oil-in-water systems that match intestinal fat emulsification requirements.

Feed technologists achieve better gut lipid absorption by selecting emulsifiers with appropriate HLB values, which optimise emulsification processes in the digestive system.

The supplementation of exogenous emulsifiers enhances bile salt function primarily in situations where bile production is insufficient or lipid intake exceeds normal levels, basically the current reality.

Field-based validation: evaluating a novel bio-emulsifier under commercial broiler conditions

A new trial approach was created to measure the practical effectiveness of a contemporary bio-emulsifier in real-world poultry farming environments. A research design incorporated mini-pen systems that operated within commercial broiler house conditions to achieve high replication numbers and authentic environmental exposure. Running tests in the animals' "natural" habitat produces genuine life data instead of depending on scientific clean, low-pressure settings.

The research involved 720 male Ross 308 chicks who received three different diets during five weeks of study.



The study involved three dietary groups that included:

- Positive Control (PC): Baseline energy diet with standard metabolisable energy (ME)
- Negative Control (NC): Reduced ME diet, 75 kcal/kg below the PC
- NC + Emulsifier: Reduced ME diet supplemented with the tested bio-emulsifier (Maxilys Plus)

Performance metrics revealed that ME restriction in the NC group caused poor growth performance with lighter final body weights and higher feed conversion ratios. This was to be expected. The addition of the bio-emulsifier to the NC diet resulted in superior performance results, which were similar to those of the PC group. The emulsifier treatment resulted in better carcass quality through reduced abdominal fat deposition along with higher breast muscle yield, which reflects improved nutrient utilisation and meat quality. Welfare indicators showed better results in the emulsifier group, indicating that this technology could bring additional benefits to gut and litter conditions.

Economic evaluation and practical takeaways

The financial analysis revealed that the bio-emulsifier addition resulted in substantial economic benefits. The positive return on investment (ROI) occurred because feed costs decreased while animal performance either remained stable or improved.

The study confirms that technologically advanced bio-emulsifiers with specific physicochemical properties that have proven effectiveness in animal trials function as important nutritional tools for modern poultry production. The additives improve energy utilisation and produce better carcass quality while delivering improved welfare outcomes and enhanced economic performance.

Conclusion

Poultry production can benefit from precision-formulated emulsifiers as a solution to fat digestion challenges while promoting sustainability and profitability. The additives work together with the bird natural emulsification processes to improve dietary lipid availability which creates new opportunities for flexible and cost-effective feed formulation. The integration of emerging research findings will become essential to preserve performance standards in an industry that faces mounting economic and environmental pressures. ■