Decoding the Secret Language of Milk

Eile van der Gaast, LifeStart Marketing Manager at Trouw Nutrition
www.trouwnutrition.com

When we feed milk to a calf, we usually think in terms of calories and nutrients supporting growth. While that is true, LifeStart research has shown that something much deeper takes place within the calf's gut.

Cow milk serves as a communication system, delivering biochemical signals that guide development and help the calf transition through its earliest stages of life. These signals do more than support short-term growth.

They shape the physiological pathways that influence how that animal performs months and years later. We've spent decades asking what calves need to grow. We are now starting to understand what milk is telling them to become.

How milk "speaks" to the calf

The "secret language" of milk is written in its complex fatty acid profile. Research has shown that specific fats in cow's milk act as signalling agents that regulate the development of the gastrointestinal (GI) tract.



At birth, the calf's GI tract is highly permeable. That temporary permeability allows colostrum immunoglobulins to pass through the intestinal lining and into the circulation, establishing passive immunity.

As the cow transitions from colostrum to milk, the composition of fatty acids changes. These fatty acids then signal the closure of the intestinal lining, forming a functional barrier against pathogens. Fatty acid messengers are vital to the calf's developmental and metabolic processes.

This shift gives calves a stronger start and accelerates maturation. These early fatty acids are not simply nutrition. They are programming the calf's future production in the herd.

Where conventional milk replacers fall short – a biological mismatch

If milk is a communication system, then the question becomes: what happens when those signals are incomplete? Calf milk replacers (CMRs) are formulated with a different priority. The focus has traditionally been on delivering cost-effective calories and nutrients to carry calves into maturity.

This mindset has often led to formulations that rely more heavily on lactose and vegetable fats, resulting in lower total fat levels than in whole milk. While whole milk contains roughly 30% fat, many CMRs are formulated with only 16–20% fat. From a nutritional standpoint, these formulations can support growth. From a biological standpoint, however, they do not fully replicate the signalling profile of natural milk and therefore do not support lifetime performance.



One of the most important differences lies in the presence of short-chain fatty acids, particularly butyric (C4:0) and caproic (C6:0) acids. These fatty acids, naturally present in cow's milk, play a role in gut development, digestive function, and metabolic regulation.

However, they are largely absent from conventional milk replacers or supplied in forms that do not mimic how they are delivered in milk. The absence of these signalling fatty acids means that development can still be supported, but it is rarely optimised. This creates a biological mismatch between how calves are designed to develop and how they are being fed.

Shifting from feeding to metabolic programming

As understanding of early-life development evolves, so too does the way we think about calf nutrition. Rather than optimising milk replacers solely for caloric and nutritional intake, the focus is shifting towards delivering the right biological signals.

One approach has been the inclusion of short-chain fatty acids in forms that more closely resemble their natural state in milk fat.

Rather than supplying butyrate or caproate as salts, which differ structurally from milk triglycerides, newer formulations incorporate compounds such as tributyrin (a source of C4:0) and tricaproin (a source of C6:0). These triglyceride forms more closely mimic how these fatty acids are delivered in whole milk and how they are digested by the calf.

Research indicates that delivering these fatty acids in triglyceride form can enhance gut development, improve digestive function, and influence systemic metabolism. They set the trajectory for how the calf's metabolism functions long after the milk-feeding phase has ended. When signalling pathways in the gut are properly activated, they influence nutrient utilisation, immune function, and energy metabolism in ways that persist into later life.

What happens when signals are right

When these biological signals are delivered at the right time and in the right form, calves show stronger growth, improved health, and more consistent development.

Studies evaluating milk replacers enriched with tributyrin and tricaproin have shown:

- Reduced need for medical interventions, including lower risk of respiratory treatments and antibiotic use
- Fewer total days requiring treatment, indicating improved resilience during early life
- Increased voluntary intake of milk replacer and starter feed, particularly around weaning
- Higher average daily gain and improved early growth trajectories

These outcomes reflect a more stable developmental environment. Calves that maintain consistent intake and experience fewer health disruptions are better able to direct nutrients towards growth rather than recovery.

The role of signalling becomes even clearer when energy intake is held constant. In studies where calves were fed high-fat milk replacers with equivalent energy levels, those receiving formulations enriched with specific fatty acids, such as tributyrin and tricaproin, consistently outperformed controls. These calves consumed more starter feed, achieved higher average daily gain during both pre-weaning and weaning, and maintained more consistent growth patterns.Because both groups received comparable energy, these differences cannot be explained by caloric intake alone. Instead, they highlight the importance of fatty acid composition and structure in regulating appetite, gut development, and metabolic function.

These early differences extend beyond the calf phase. A meta-analysis of 12 studies found that for every additional kilogram of average daily gain achieved during the pre-weaning phase, dairy heifers produced approximately 1,500 kilograms more milk in their first lactation. This is the foundation of metabolic programming. Early-life nutritional signals shape how efficiently the animal utilises nutrients, responds to stress, and performs over its lifetime.

For dairy producers, the implications are practical. Calves that grow more consistently and remain healthier reach key milestones earlier, including breeding weight, reducing rearing costs by €40 to €60 per animal.

At the same time, fewer health interventions and improved resilience reduce management inputs while increasing overall herd productivity.

Milk speaks a language to the calf's gastrointestinal tract that we are only beginning to translate. The next advance in calf nutrition will not be measured solely in grams of protein or fat, but in how precisely we can replicate the biological signals that nature designed. ■