The hatchery's role in unlocking genetic potential:

Why Incubation Management Matters More than Ever in Modern Poultry Genetics

Bert Peters, Global Hatchery Specialist, Hendrix Genetics
layinghens.hendrix-genetics.com



The poultry industry often credits genetic progress for improvements in egg production, feed efficiency, persistency and liveability. These gains are real, but genetics only creates value when birds can fully express their potential under commercial conditions.

The hatchery plays a critical role in that process. A modern layer expected to remain productive beyond 100 weeks of age must arrive on the rearing farm with a well-developed skeleton, cardiovascular system, gastrointestinal tract and immune system. Much of this development takes place during incubation. While a chick may survive suboptimal incubation conditions, survival should not be confused with optimal development. Decisions made during incubation can influence biological development long before differences become visible in flock performance.

Modern laying hens are not the same birds they were 20 years ago

Today's commercial laying hens produce significantly more egg mass, remain productive for longer periods and are expected to maintain performance under increasingly demanding production conditions. As a result, developmental mistakes during incubation have become more costly than ever before.

Continued on page 12 ▷

▷ *Continued from page 11*

Historically, incubation programmes were largely optimised around hatchability. While hatchability remains an important metric, modern hatcheries should increasingly focus on producing chicks capable of expressing their genetic potential throughout a laying cycle that may exceed 100 weeks.

This shift changes the question we ask. Instead of focusing solely on how many chicks hatched, we should be asking how well those chicks were developed. The difference may seem subtle, but it fundamentally changes how we evaluate hatchery performance.

Embryo temperature remains the most important biological parameter

Among all incubation parameters, embryo temperature remains the single most important driver of embryonic development. Most hatcheries closely monitor machine temperatures, yet embryos respond to their own temperature rather than to incubator set points.

This distinction becomes increasingly important during the second half of incubation, when metabolic heat production rises rapidly. Excessive embryo temperatures are commonly associated with:

- Accelerated embryonic development
- Earlier hatch
- Increased dehydration
- Poorer yolk utilisation
- Reduced chick quality

Conversely, low embryo temperatures can result in:

- Delayed hatch
- Increased late embryonic mortality
- Reduced physiological maturity at hatch

Managing embryo temperature becomes even more challenging because not all embryos produce the same amount of metabolic heat. Factors such as:

- Breeder age
- Egg size
- Genetic strain
- Egg storage duration
- Incubation uniformity

all influence embryonic metabolism and heat production. Consequently, identical machine settings do not necessarily create identical developmental conditions. For this reason, successful hatcheries increasingly focus on embryo response rather than machine settings alone. The question is no longer, "What temperature is the machine running?" but rather, "What temperature is the embryo experiencing?"

Hatch window is often a symptom, not the real problem

This brings us to another parameter that receives considerable attention in modern hatcheries: hatch window.

Many hatcheries work hard to reduce hatch window length, but the hatch window itself is rarely the root cause of poor performance. More often, it is a visible symptom of variation introduced much earlier in the process.

A wide hatch window is commonly associated with variation in:

- Breeder flock uniformity
- Breeder flock age
- Egg size
- Storage conditions
- Prewarming procedures
- Temperature distribution during incubation

By the time chicks begin hatching, the underlying variation has already occurred. From a biological standpoint, a prolonged hatch window creates variation in physiological age. The earliest chicks may spend 24 to 36 hours waiting before placement, while the latest chicks are still completing embryonic development. At processing, both are classified as day-old chicks. Biologically, however, they are at very different stages of development.

This variation does not disappear after placement. In many cases, it follows the flock throughout rearing and can still be observed later in production through differences in body weight, uniformity and performance.

Uniform incubation produces uniform flocks

The importance of hatch window becomes even clearer when considering flock uniformity.

Throughout the poultry industry, considerable effort is invested in achieving uniform pullet flocks. However, uniformity does not start in the rearing house. It starts in the incubator.

When embryos experience differences in temperature, oxygen availability or moisture loss, developmental rates begin to diverge. These differences can influence:

- Hatch timing
- Chick body weight
- Residual yolk absorption
- Intestinal development
- Early feed intake

Over time, these relatively small differences can develop into measurable variation within the flock.

Many hatcheries have invested heavily in modern equipment while assuming that every position within a machine experiences identical incubation conditions. Field audits repeatedly show that this is often not the case. Temperature mapping, airflow distribution measurements and eggshell temperature assessments frequently reveal variation that is not apparent from machine controllers alone.

As a result, leading hatcheries are increasingly focused on reducing variation rather than simply maximising averages. Uniform incubation creates uniform chicks, and uniform chicks create uniform flocks.

Ventilation influences much more than hatchability

When ventilation is discussed, the conversation often focuses on oxygen supply and carbon dioxide removal. In reality, its influence is much broader.

Airflow distribution directly affects:

- Heat removal
- Embryo temperature
- Moisture loss
- Hatch synchrony
- Navel quality

In many investigations where hatchability differs between locations within the same machine, airflow distribution ultimately proves to be one of the underlying causes.

The embryo does not experience average machine conditions. It experiences its local environment. Small differences in airflow may seem insignificant over a few hours, but over a 21-day incubation period they can create meaningful differences in embryonic development.

For that reason, ventilation should be viewed as a developmental tool, not simply an engineering requirement.

The chick's first 24 hours remain underestimated

The biological significance of incubation does not end at hatch.

In many operations, attention shifts away from the chick once it leaves the hatcher. Yet some of the most important developmental processes are still taking place during the first hours after emergence.

During this period:

- Residual yolk is absorbed
- Digestive function accelerates
- Thermoregulation develops
- Immune maturation continues

The industry has become increasingly aware that chick quality is not defined at hatch, but at placement.

The objective should therefore be to minimise the time between emergence and access to feed and water while maintaining chick comfort throughout processing, holding and transport. As production cycles continue to become longer, the economic consequences of early developmental setbacks become increasingly significant.

A small disadvantage created during the first day of life may still be measurable more than 100 weeks later.

The future of incubation is developmental biology

The hatcheries that consistently outperform others are no longer focused exclusively on hatchability percentages. Increasingly, they monitor biological indicators such as:

- Embryo temperature
- Hatch window
- Chick activity
- Yolk utilisation
- Navel quality
- Chick uniformity

In other words, they evaluate how incubation influenced development, not simply whether incubation was completed successfully. This is where genetics and hatchery management increasingly converge. Modern breeding programmes continue to deliver birds with greater potential for egg production, feed efficiency, liveability and lifetime performance. However, genetic progress only creates value when embryos are allowed to develop under conditions that support the expression of that potential.

As production cycles become longer and birds become more efficient, the cost of developmental mistakes during incubation increases. The objective must be to consistently produce physiologically mature, robust and uniform chicks that are fully prepared for the demands of modern production systems. The next step is to deliver these chicks successfully and in good condition to the rearing farm! ■

Geneticists can build the blueprint for a 600-egg hen, but it is the hatchery that determines whether that blueprint becomes a reality.

